2.

1


Denver Wildlife Research Center, U.S.Department of Interior August 1969

## Rate of Hydrolysis of Zinc Phosphide

- Eight analyses were run, using solutions ranging in pH from about 0.2 through about 14.
- 2. In each analysis, 100 ml. of water (adjusted to a given pH with buffers, acid, or base) was added to a reaction flask containing known amounts of Zn<sub>3</sub>P<sub>2</sub>. The flask was maintained at 50°C. and a continuous N<sub>2</sub> sweep was maintained for 1 hour.
- 3. All analyses were run by the method set forth in the Federal Specification for  $\text{Zn}_3P_2$  except for substituting various buffered pH solutions for the sulfuric acid.
- 4. Results were as follows (also see graph titled "Hydrolysis Rate"):

	Solution used	рн	Recovery
1.	10.0% v/v sulfuric acid in H20 ca.	0.2	97•59
2.	Coleman Buffer, certified	4.0	51.73
3.	n n	6.0	5.98
4.	Boiled, distilled H20	7.0 <del>*</del>	0.00
5.	Coleman Buffer, certified	8.0	1.24
6.	и и	10.0	1.12
7•	(E	12.4 andbook	16.40
8		14	70.02

\* At a pH of 7 (distilled H<sub>2</sub>O only) no reduction of the potassium permanganate could be observed; therefore, in this case the reaction time was extended to 2 hours.



Comments by W. H. Robison, Chemist

## Conclusions reached during this investigation:

- 1. There are several factors governing the recovery of PH<sub>3</sub> generated by hydrolysis of Zn<sub>3</sub>P<sub>2</sub> in the presence of sample material. These include:
  - A. pH of reaction flask contents.
  - B. Oxidation of PH<sub>3</sub> to non-volatile components -probably phosphates.
  - C. Irreversible adsorption of PH<sub>3</sub> or its equivalent as phosphorus-containing compounds by the sample material.
- 2. The method will detect  $PH_3$  at the 0.005 ppm level and givereliable quantitative results at the 0.05 ppm level or above.

T BEST DOCUMENT AVAILABLE

a: Jackson Rowen 16-27-83

## DATA EVALUATION RECEDEN

Yackson Cowen 10-27-53

FEEF

**コ**フフ 10450030

DATA EVALUATION RECORD

PAGE

CASE GS0026

ZINC PHOSPHIDE

PM 400 01/28/50

CHE# 088601

Zine phosphide

RRANCH EFE DISC 30 TOPIC 05:015

FORMULATION OF - ACTIVE INGREDIENT

FICHE/MASTER ID 00005926

U.S. Fish and mildlife Service, Denver mildlife Pesearch Center (1969) Rate of Hydrolysis of Zinc phosphide. (Unpublished study including comments by w.H. Robinson, received Sep 25, 1969 under OFOR90; submitted by Maweil, Dept. of Agriculture, homolulu, Hawaii; CDL:093187-K)

SURST. CLASS = 5.

DIMECT RYN TIME & 0.25 (MH) STAPT-DATE 3/23/8END DATE 2/13/82

TITLE:

see allahuel sheet-

SIGNATURES

LOC/TEL:

Edward BB atter 011E1 2/23/8=

APPROVED BY:

TITLE

DRG: LOC/TEL:

SIGNATURE:

DATES

CONCLUSIONS

This submission with data showing that
Accords solutions of zinc phosphide are Stable only at pt/ 20 is per imont to the Registration standard, since the product chemistry quidelines require data in agreous stability of each

## DATA EVALUATION RECORD

Vackson Rowen 10-27-83

P=EF

TOMSCOZO

DATA EVALUATION RECORD

PAGE

CASE GS0026

ZINC PHOSPHIDE

**ユフフ ・・・・** 

PM 400 01/28/80

CHEM 088601

Zine phosphide

RRANCH EFE

DISC 30 TOPIC 051015

FORMULATION OF - ACTIVE INGREDIENT

FICHE/HASTER ID 00005926

CUNTENT CAT 02

U.S. Fish and wildlife Sarvice, Denver Wildlife Pesearch Center (1969) Rate of Hydrolysis of Zinc phosphide. (Unpublished study including comments by W.H. Robinson, received Sep 25, 1969 under OFOR90; submitted by Hawaii, Dept. of Agriculture, Honolulu, Hawaii; CDL:093187-K)

SURST. CLASS # 5.

DIRECT RYN TIME = 0.25 (MH) STAPT-DATE 2/23/FLEND DATE 2/13/82

PEVIENED BY:

TITLE: OkG:

LOC/TEL:

See aliable sheet

SIGNATURES

Edward B-B cutter 011E1 2/23/82

APPROVED BY:

TITLE

ORG:

LOCATEL:

SIGNATURE:

DATES

CONCLUSIONS

This submission with data showing that?
Accords solutions of zine phosphide are Stable only at pt/ 20 is pictiment to the Registration standard, since the product chemistry quitelines require data en aquious stability of each